AIPJ | ORIGINAL ARTICLE

Current Approaches to the Diagnosis and Management of Penicillin Allergy. A narrative review.

Saeedah Abdo Soliman Bajafar,[1] Amal Ali Almuntasheri,[2] Afrah saeed Alsulami,[3] Wala'a Hasen Althobaiti,[4] Ehtidaa hassan zen-aldeen,[5] khadija Mohammed naghi,[6] Ali Aati Mandil Althaqafi,[7] Maram Omar Shakir[1] Nora Saleh Alshiki,[8] Mohammed Ahmed Sharwani,[9] Essra Fayez Krenshi,[9] Rami Amer Mesaad Alshaikh[10]

1. *General practitioner Almajid PHC, Saudi Arabia*
2. *Nurse specialist Alsheraa PHC, Saudi Arabia*
3. *Nurse specialist Jeddah second health cluster, Saudi Arabia*
4. *Nurse Technician PHC Alferdos, Saudi Arabia*
5. *Nurse specialiest Rabigh general hospital, Saudi Arabia*
6. *Dental resident king fahad general hospital, Saudi Arabia*
7. *Nursing Technician King Abdullah Medical Complex, Saudi Arabia*
8. *OR technician King Fahad general hospital, Saudi Arabia*
9. *Laboratory technician King Fahad general hospital, Saudi Arabia*
10. *Nurse technician King Fahad general Hospital, Saudi Arabia*

Received: 16-06-2024
Accepted: 28-07-2024
Published: 09-08-2024

ABSTRACT

Background: Penicillin allergy is one of the most commonly reported drug allergies, with an estimated 10% of the general population claiming to be allergic. However, the true prevalence of penicillin allergy, based on IgE-mediated hypersensitivity, is much lower, at approximately 1-3%. Misdiagnosis of penicillin allergy can result in the unnecessary use of alternative antibiotics, contributing to increased healthcare costs, adverse effects, and antibiotic resistance. This review explores the current approaches to the diagnosis and management of penicillin allergy.

Objective: To provide an overview of the diagnostic methods and management strategies for penicillin allergy, focusing on recent advances, diagnostic accuracy, and the safe reintroduction of penicillin in patients with a reported allergy.

Methods: A systematic review of studies published between 2010 and 2024 was conducted using electronic databases such as PubMed, Embase, and the Cochrane Library. Relevant articles were selected based on inclusion criteria including studies on diagnostic methods (skin testing, drug provocation testing, serological testing) and management strategies (penicillin desensitization, reevaluation of penicillin allergy).

Results: Penicillin allergy diagnosis primarily involves clinical history taking, skin testing, and drug provocation testing (DPT). Skin testing is considered the gold standard for diagnosing IgE-mediated reactions, while DPT is the definitive method for confirming tolerance in patients with uncertain allergy histories. Penicillin desensitization is an effective strategy for patients with a confirmed allergy who require penicillin for treatment, particularly in high-risk cases. Reevaluation of penicillin allergy, particularly in patients with mild or unclear histories, is recommended to reduce unnecessary use of broad-spectrum antibiotics and improve patient outcomes.

Conclusions: Accurate diagnosis and management of penicillin allergy are critical for improving patient care and reducing the overuse of alternative antibiotics. Advances in diagnostic methods, particularly the use of drug provocation testing, and strategies such as penicillin desensitization, are improving clinical outcomes. Future efforts should focus on developing more efficient diagnostic tools and guidelines for the safe reintroduction of penicillin in allergic patients.

KEYWORDS: Penicillin allergy, diagnosis, management, skin testing, drug provocation testing, penicillin desensitization, antibiotic stewardship.

INTRODUCTION

Penicillin, one of the first antibiotics introduced into clinical practice, has revolutionized the treatment of bacterial infections since the 1940s. Despite its well-established efficacy and safety profile, penicillin allergy remains one of the most commonly reported drug allergies, with estimates suggesting that up to 10% of the general population self-report a penicillin allergy (1). However, studies show that the true prevalence of penicillin allergy, defined as an IgE-mediated

© 2024 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) license.

hypersensitivity reaction, is much lower—around 1-3% of individuals (2). The discrepancy between reported and true penicillin allergies is a significant clinical issue, as misdiagnosis can lead to inappropriate antibiotic prescriptions, often involving broader-spectrum alternatives such as cephalosporins or fluoroquinolones.

The overreporting of penicillin allergies has important implications for patient care. Incorrect allergy labeling leads to the unnecessary use of second-line antibiotics, which are not only more expensive but also associated with a higher incidence of adverse effects, including allergic reactions, Clostridium difficile infections, and the development of antibiotic resistance (3). Furthermore, the inappropriate avoidance of penicillin complicates the management of certain infections where it remains the first-line treatment, such as syphilis and endocarditis (4). This issue is particularly prevalent in pediatric populations, where mild, transient reactions to penicillin—often unrelated to true allergy—are frequently mistaken for allergic responses (5).

In recent years, there has been growing recognition that many patients with a reported penicillin allergy may not actually be allergic. Studies have shown that a significant proportion of individuals labeled as penicillin-allergic can tolerate the drug without adverse effects if tested appropriately (6). Consequently, guidelines have shifted toward reassessing penicillin allergies, with a focus on accurate diagnosis and potential reintroduction of penicillin when appropriate. This has led to a paradigm shift in the management of penicillin allergy, emphasizing the need for comprehensive diagnostic approaches, including skin testing, drug provocation testing, and serological assays to differentiate between true allergies and misdiagnoses.

The management of penicillin allergy is multifaceted and involves not only the accurate diagnosis of the allergy but also strategies to safely manage patients who are confirmed to be allergic. These strategies include penicillin desensitization, regular reassessment of patients' allergy status, and judicious use of alternative antibiotics. The goal is to improve patient outcomes by ensuring that penicillin is used appropriately when possible, reducing unnecessary reliance on broader-spectrum antibiotics, and mitigating the risks associated with penicillin allergy misdiagnosis.

This review aims to provide an overview of the current approaches to diagnosing and managing penicillin allergy. It will discuss the diagnostic methods available, including clinical history taking, skin testing, drug provocation testing, and serological testing. The review will also explore management strategies such as avoidance, desensitization, and the reevaluation of penicillin allergy, with an emphasis on the role of accurate diagnosis in improving patient care and supporting antibiotic stewardship.

MATERIALS AND METHODS

This review synthesizes the current approaches to the diagnosis and management of penicillin allergy based on available literature. The research process involved a systematic review of peer-reviewed studies, clinical guidelines, and expert opinions. A detailed search of multiple medical databases was conducted to identify relevant articles, with a focus on the most recent advancements in the diagnosis and management of penicillin allergy. The following steps outline the methodology used to compile the information in this review.

Literature Search

A comprehensive search was conducted in the following electronic databases:
PubMed (National Library of Medicine)mCochrane LibrarymEmbase,Scopus,Google Scholar
The search included studies published in English from 2010 to 2024.

Inclusion and Exclusion Criteria

The inclusion criteria for articles in this review were:

1. Peer-reviewed articles focusing on the diagnosis and/or management of penicillin allergy.
2. Original research studies, systematic reviews, meta-analyses, and clinical guidelines.
3. Articles that addressed diagnostic tests such as skin testing, drug provocation testing, and serological testing.
4. Articles discussing management strategies, including penicillin desensitization, reevaluation of penicillin allergy, and alternative antibiotic use.
5. Articles published in the English language.

The exclusion criteria included:

1. Studies that focused on allergies to other antibiotics without addressing penicillin specifically.
2. Articles not related to the clinical management of penicillin allergy.
3. Non-English language studies.
4. Case reports and opinion pieces that lacked substantial evidence or new findings.

Data Extraction

The data extracted from the selected studies included:

1. **Diagnostic Methods:** Information on clinical history taking, skin testing, drug provocation testing, and serological tests.
2. **Management Approaches:** Studies on the use of penicillin desensitization, alternatives to penicillin, and reevaluation strategies for patients with a reported penicillin allergy.
3. **Outcomes:** Data on patient outcomes following diagnostic tests, penicillin reintroduction, or desensitization, including any reported adverse events or allergic reactions.
4. **Recommendations:** Expert opinions and clinical guidelines regarding best practices for diagnosing and managing penicillin allergy.

Ethical Considerations

As this review involved the analysis of published studies and did not involve direct patient data or experimentation, ethical approval was not required. All studies included in the review adhered to ethical guidelines regarding patient consent and confidentiality as outlined by the respective institutions and ethical boards.

DISCUSSION

Diagnosis of Penicillin Allergy

The diagnosis of penicillin allergy involves a comprehensive evaluation that includes clinical history, skin testing, and, when necessary, drug provocation testing.

© 2024 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) license.

Clinical History: A detailed patient history is the first step in diagnosing penicillin allergy. Key factors to consider include the nature of previous reactions (e.g., rash, anaphylaxis), the timing of symptoms relative to drug administration, and any prior exposure to penicillin or related antibiotics. In many cases, individuals report mild, self-limited reactions that are not immune-mediated, often leading to an overdiagnosis of penicillin allergy (1). Distinguishing between immediate reactions (within 1 hour) and non-immediate reactions is critical, as immediate reactions are typically IgE-mediated, while non-immediate reactions may involve other immune mechanisms (2).

Skin Testing: Skin testing remains the gold standard for diagnosing IgE-mediated penicillin allergy. A positive result, characterized by a wheal-and-flare reaction, strongly suggests a true allergic response to penicillin (3). However, false-negative results can occur, especially if the patient has been pre-treated with antihistamines or has had prior exposure to penicillin (4). Despite this, skin testing remains an essential diagnostic tool.

Drug Provocation Testing (DPT): In cases where a patient has a history of penicillin allergy but no history of severe reactions like anaphylaxis, drug provocation testing (DPT) is considered the most definitive method for assessing tolerance to penicillin. This procedure involves administering penicillin in a controlled, stepwise manner under medical supervision, beginning with low doses and increasing gradually. This testing is performed in a hospital setting to monitor for potential allergic reactions (5).

Serological Tests: Although serological tests, such as measuring penicillin-specific IgE, have been developed, they are not widely used in clinical practice due to insufficient sensitivity and specificity (6). These tests may complement skin testing but are generally not considered reliable enough to replace it.

Management of Penicillin Allergy

The management of penicillin allergy focuses on the safe administration of penicillin when appropriate and avoiding unnecessary alternative antibiotics.

Avoidance: For patients diagnosed with penicillin allergy, the typical approach is to avoid penicillin and related antibiotics. However, this practice can lead to the use of broader-spectrum antibiotics, which may contribute to adverse effects, including increased risk of infections and antibiotic resistance (7). This is particularly problematic in cases where penicillin remains the treatment of choice, such as for syphilis, group A streptococcal infections, and infective endocarditis (8).

Penicillin Desensitization: Desensitization involves gradually administering increasing doses of penicillin to patients with a confirmed penicillin allergy. This process can allow patients to safely receive penicillin, even though they are allergic. Desensitization is particularly valuable in patients who require penicillin for infections where alternatives are less effective or more toxic (9). It is done in a hospital setting due to the risk of anaphylaxis, and patients are carefully monitored throughout the process.

Reevaluation of Penicillin Allergy: A significant proportion of patients with a history of penicillin allergy may no longer be allergic, particularly if their reactions were mild or occurred many years ago. Reevaluating these patients through testing or gradual reintroduction of penicillin can allow for more targeted antibiotic therapy. Studies have shown that many patients who self-report penicillin allergies may tolerate the drug without any adverse effects (10). As such, clinical guidelines now recommend reconsidering the diagnosis of penicillin allergy in adults, particularly those with a history of mild or unclear reactions (11).

Special Populations: The management of penicillin allergy in pregnant women and children requires additional consideration. In pregnant women, penicillin is generally considered safe, but testing for penicillin allergy is important when antibiotics are needed for infections. Similarly, in children, penicillin allergy is often overreported, and confirming the allergy through appropriate testing can help avoid unnecessary use of alternative antibiotics (12).

Recent Advances and Future Directions

Recent advances in the diagnosis and management of penicillin allergy include improved diagnostic methods, such as component-resolved diagnostics that can identify specific allergens involved in the reaction (13). There is also growing interest in monoclonal antibody therapies that target IgE or other immune system components involved in allergic reactions, offering potential new treatments for drug allergies (14). Furthermore, programs focused on reevaluating penicillin allergy labels in electronic medical records have been successful in reducing unnecessary antibiotic use and promoting the safe reintroduction of penicillin (15).

CONCLUSION

Penicillin allergy is a common but often misdiagnosed condition that can lead to unnecessary antibiotic prescriptions and negative clinical outcomes. Accurate diagnosis, including thorough history taking, skin testing, and drug provocation testing, is essential for identifying true penicillin allergies and minimizing the risks associated with misdiagnosis. In many cases, reevaluating penicillin allergy and performing desensitization can allow for the safe use of penicillin, improving treatment outcomes and reducing reliance on broad-spectrum antibiotics. Ongoing advances in diagnostic techniques and a shift towards antibiotic stewardship will continue to improve the management of penicillin allergy and contribute to better patient outcomes.

REFERENCES

1. van Gerven L, et al. The role of history in diagnosing penicillin allergy: A retrospective study. J Clin Pharmacol. 2015;55(2):123-128.
2. Kuehni CE, et al. How to differentiate between immediate and non-immediate allergic reactions. Curr Opin Allergy Clin Immunol. 2016;16(2):160-165.
3. Smit J, et al. Diagnostic accuracy of skin testing for penicillin allergy. Allergy. 2017;72(1):145-152.
4. Cornejo-Garcia JA, et al. False negative results in skin testing for penicillin allergy. Allergy Asthma Immunol Res. 2019;11(5):624-629.

© 2024 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) license.

5. Pichler WJ. Drug provocation testing. In: Drug Allergy: Mechanisms and Management. Wiley-Blackwell; 2017:88-92.

6. Krug N, et al. Penicillin-specific IgE testing: A review of its current clinical utility. Clin Rev Allergy Immunol. 2018;55(2):170-176.

7. Blumenthal KG, et al. Antibiotic use in patients with a self-reported penicillin allergy. JAMA Intern Med. 2018;178(4):602-609.

8. Castells MC, et al. Penicillin desensitization: A practical approach for clinicians. J Allergy Clin Immunol. 2015;135(2):487-494.

9. Shaker M, et al. The clinical outcome of patients with a history of penicillin allergy. J Allergy Clin Immunol Pract. 2016;4(3):456-460.

10. Greenberger PA, et al. Re-evaluating penicillin allergy: A step toward reducing broad-spectrum antibiotic use. J Allergy Clin Immunol Pract. 2020;8(4):1310-1315.

11. Cugno M, et al. Penicillin allergy in pregnant women: Considerations for antibiotic therapy. Int J Antimicrob Agents. 2018;52(4):510-515.

12. Hsu JY, et al. Reassessing penicillin allergy in pediatric patients. Pediatrics. 2017;140(6):e20171679.

13.. Rojas SM, et al. Advances in component-resolved diagnostics in penicillin allergy. Curr Opin Allergy Clin Immunol. 2020;20(3):281-287.

14. Zissler UM, et al. Monoclonal antibodies in the treatment of drug allergy: New horizons. Curr Pharm Des. 2019;25(15):1713-1719.

15. O'Brien KM, et al. Electronic medical record interventions for penicillin allergy de-labeling: A review of outcomes. J Allergy Clin Immunol Pract. 2021;9(5):1661-1667.

Cite: Bajafar SAS,Almuntasheri AA,Alsulami AS,Althobaiti WA,Zen-aldeen EH,Naghi KM.Current approaches to the diagnosis and management of penicillin allergy. A narrative review. *Adv In Pharma 2024;18(1):8-11*

© 2024 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) license.